

Facility Quality Measure/Indicator Report

Facility Name XXXXXXXXXXXXXXXX
City/State DALLAS, TX
Provider Number XXXXXX
Login/Facility ID XXXXXXXXXXXX
Data was calculated on 07/16/2007

Run Date 07/20/07 08:48:08
Report Period 01/01/07 - 06/30/07
Comparison Group 11/01/06 - 04/30/07
Report Version Number 1.07

Measure ID	Domain/Measure Description	Facility				Comparison Group		
		Num	Denom	Observed Percent	Adjusted Percent	State Average	National Average	State Percentile
Chronic Care Measures								
<u>Accidents</u>								
1.1	Incidence of new fractures	1	114	0.9%	-	2.0%	2.0%	44
1.2	Prevalence of falls	14	118	11.9%	-	10.9%	13.0%	61
<u>Behavior/Emotional Patterns</u>								
2.1	Residents who have become more depressed or anxious	7	113	6.2%	-	9.3%	15.5%	41
2.2	Prevalence of behavior symptoms affecting others: Overall	27	115	23.5%	-	14.1%	17.7%	84
2.2-HI	Prevalence of behavior symptoms affecting others: High risk	26	106	24.5%	-	16.3%	20.5%	81
2.2-LO	Prevalence of behavior symptoms affecting others: Low risk	1	9	11.1%	-	6.0%	7.5%	81
2.3	Prevalence of symptoms of depression without antidepressant therapy	3	115	2.6%	-	2.3%	4.9%	68
<u>Clinical Management</u>								
3.1	Use of 9 or more different medications	65	118	55.1%	-	66.1%	65.8%	17
<u>Cognitive Patterns</u>								
4.1	Incidence of cognitive impairment	0	12	0.0%	-	13.2%	12.8%	0
<u>Elimination/Incontinence</u>								
5.1	Low-risk residents who lost control of their bowels or bladder	29	72	40.3%	-	44.1%	48.3%	38
5.2	Residents who have/had a catheter inserted and left in their bladder	6	118	5.1%	5.0%	7.0%	7.5%	43
5.3	Prevalence of occasional or frequent bladder or bowel incontinence without a toileting plan	7	15	46.7%	-	71.2%	45.6%	23
5.4	Prevalence of fecal impaction	0	118	0.0%	-	0.1%	0.1%	0
<u>Infection Control</u>								
6.1	Residents with a urinary tract infection	7	118	5.9%	-	7.8%	9.7%	42
<u>Nutrition/Eating</u>								
7.1	Residents who lose too much weight	6	109	5.5%	-	8.0%	10.3%	38
7.2	Prevalence of tube feeding	13	118	11.0%	-	8.0%	6.8%	75
7.3	Prevalence of dehydration	0	118	0.0%	-	0.1%	0.3%	0
<u>Pain Management</u>								
8.1	Residents who have moderate to severe pain	3	118	2.5%	2.1%	5.8%	5.6%	31
<u>Physical Functioning</u>								
9.1	Residents whose need for help with daily activities has increased	9	93	9.7%	-	16.8%	17.6%	22
9.2	Residents who spend most of their time in bed or in a chair	2	115	1.7%	-	6.9%	5.3%	23
9.3	Residents whose ability to move in and around their room got worse	4	80	5.0%	4.8%	13.6%	15.9%	15
9.4	Incidence of decline in ROM	2	116	1.7%	-	6.6%	8.1%	20

Note: Dashes represent a value that could not be computed

Facility Quality Measure/Indicator Report

Facility Name XXXXXXXXXXXXXXXX
City/State DALLAS, TX
Provider Number XXXXXX
Login/Facility ID XXXXXXXXXXXX
Data was calculated on 07/16/2007

Run Date 07/20/07 08:48:08
Report Period 01/01/07 - 06/30/07
Comparison Group 11/01/06 - 04/30/07
Report Version Number 1.07

Measure ID	Domain/Measure Description	Facility				Comparison Group		
		Num	Denom	Observed Percent	Adjusted Percent	State Average	National Average	State Percentile
Chronic Care Measures								
Psychotropic Drug Use								
10.1	Prevalence of antipsychotic use, in the absence of psychotic or related conditions: Overall	24	97	24.7%	-	22.2%	20.2%	63
10.1-HI	Prevalence of antipsychotic use, in the absence of psychotic or related conditions: High risk	10	19	52.6%	-	45.3%	42.9%	63
10.1-LO	Prevalence of antipsychotic use, in the absence of psychotic or related conditions: Low risk	14	76	18.4%	-	19.6%	16.9%	51
10.2	Prevalence of antianxiety/hypnotic use	10	97	10.3%	-	23.8%	20.9%	7
10.3	Prevalence of hypnotic use more than two times in last week	0	118	0.0%	-	8.0%	4.9%	0
Quality of Life								
11.1	Residents who were physically restrained	3	118	2.5%	-	4.8%	5.5%	48
11.2	Prevalence of little or no activity	3	115	2.6%	-	10.2%	7.8%	35
Skin Care								
12.1	High-risk residents with pressure ulcers	6	56	10.7%	-	13.1%	14.3%	41
12.2	Low-risk residents with pressure ulcers	0	62	0.0%	-	2.0%	2.9%	0
Post-Acute Care(PAC) Measures								
13.1	Short-stay residents with delirium	6	39	15.4%	10.2%	2.8%	2.5%	93 *
13.2	Short-stay residents who had moderate to severe pain	3	41	7.3%	-	17.3%	20.8%	27
13.3	Short-stay residents with pressure ulcers	1	33	3.0%	2.3%	15.0%	17.1%	10

Note: Dashes represent a value that could not be computed